


27. Peso total da biomassa aérea por grupo de espécies  
*Total weight of aboveground tree biomass by species group*

Folhosas/Coníferas <i>Broadleaves/Conifers</i>	Uso do solo / Biomassa aérea total <i>Land-use class / Aboveground biomass</i>			
	Santo Antão - Agroflorestal <i>Santo Antao - Agroforestry</i>		Santo Antão - Floresta <i>Santo Antao - Forest</i>	
	thousands t ( $\alpha = 0.1$ )	%	thousands t ( $\alpha = 0.1$ )	%
Folhosas <i>Broadleaves</i>	0.0	1.1	24.1 (18.1 – 30.1)	42.3
Coníferas <i>Conifers</i>	0.4	98.9	32.9 (24.2 – 41.6)	57.7
<b>Total</b>	<b>0.4</b>	<b>100.0</b>	<b>57.0 (45.7 – 68.3)</b>	<b>100.0</b>

Folhosas/Coníferas <i>Broadleaves/Conifers</i>	Uso do solo / Biomassa aérea total <i>Land-use class / Aboveground biomass</i>			
	Santo Antão - Formação florestal aberta <i>Santo Antao - Open forest</i>		Total	
	thousands t ( $\alpha = 0.1$ )	%	thousands t ( $\alpha = 0.1$ )	%
Folhosas <i>Broadleaves</i>	0.6 (0.4 – 0.8)	95.4	24.7 (18.7 – 30.7)	42.6
Coníferas <i>Conifers</i>	0.0	4.6	33.3 (24.6 – 42.0)	57.4
<b>Total</b>	<b>0.6 (0.4 – 0.9)</b>	<b>100.0</b>	<b>58.0 (46.7 – 69.3)</b>	<b>100.0</b>



27. Peso total da biomassa aérea por grupo de espécies