


28. Peso médio de biomassa aérea por hectare e por grupo de espécies  
*Mean weight of aboveground tree biomass per hectare by species group*

Folhosas/Coníferas <i>Broadleaves/Conifers</i>	Uso do solo / Biomassa aérea total <i>Land-use class / Aboveground biomass</i>			
	Santo Antão - Agroflorestal <i>Santo Antao - Agroforestry</i>		Santo Antão - Floresta <i>Santo Antao - Forest</i>	
	t/ha	( $\alpha = 0.1$ )	t/ha	( $\alpha = 0.1$ )
Folhosas <i>Broadleaves</i>	0.0	(0.0 – 0.1)	12.0	(8.9 – 15.1)
Coníferas <i>Conifers</i>	3.6	(0.0 – 10.2)	16.4	(11.1 – 21.6)
All	3.7	(0.0 – 10.3)	28.4	(22.7 – 34.0)

Folhosas/Coníferas <i>Broadleaves/Conifers</i>	Uso do solo / Biomassa aérea total <i>Land-use class / Aboveground biomass</i>			
	Santo Antão - Formação florestal aberta <i>Santo Antao - Open forest</i>		All	
	t/ha	( $\alpha = 0.1$ )	t/ha	( $\alpha = 0.1$ )
Folhosas <i>Broadleaves</i>	3.6	(2.3 – 5.0)	4.6	(3.0 – 6.2)
Coníferas <i>Conifers</i>	0.2	(0.0 – 0.5)	6.2	(3.4 – 8.9)
All	3.8	(2.5 – 5.2)	10.8	(7.8 – 13.7)



28. Peso médio de biomassa aérea por hectare e por grupo de espécies