


20. Peso total da biomassa aérea por grupo de espécies  
*Total weight of aboveground tree biomass by species group*

Folhosas/Coníferas <i>Broadleaves/Conifers</i>	Uso do solo / Biomassa aérea total <i>Land-use class / Aboveground biomass</i>					
	Santiago - Agroflorestal <i>Santiago - Agroforestry</i>			Santiago - Floresta <i>Santiago - Forest</i>		
	thousands t	( $\alpha = 0.1$ )	%	thousands t	( $\alpha = 0.1$ )	%
Folhosas <i>Broadleaves</i>	61.2	(41.1 – 81.2)	100.0	357.7	(337.7 – 377.7)	99.5
Coníferas <i>Conifers</i>	0.0		0.008	1.9	(0.0 – 3.9)	0.5
<b>Total</b>	<b>61.2</b>	<b>(41.1 – 81.2)</b>	<b>100.0</b>	<b>359.6</b>	<b>(339.5 – 379.8)</b>	<b>100.0</b>

Folhosas/Coníferas <i>Broadleaves/Conifers</i>	Uso do solo / Biomassa aérea total <i>Land-use class / Aboveground biomass</i>					
	Santiago - Formação florestal aberta <i>Santiago - Open forest</i>			Santiago - Área arbustiva <i>Santiago - Shrubland</i>		
	thousands t	( $\alpha = 0.1$ )	%	thousands t	( $\alpha = 0.1$ )	%
Folhosas <i>Broadleaves</i>	45.5	(17.7 – 73.3)	100.0	16.1	(7.4 – 24.8)	100.0
Coníferas <i>Conifers</i>	–	–	–	–	–	–
<b>Total</b>	<b>45.5</b>	<b>(17.7 – 73.3)</b>	<b>100.0</b>	<b>16.1</b>	<b>(7.4 – 24.8)</b>	<b>100.0</b>

Folhosas/Coníferas <i>Broadleaves/Conifers</i>	Uso do solo / Biomassa aérea total <i>Land-use class / Aboveground biomass</i>		
	Total		
	thousands t	( $\alpha = 0.1$ )	%
Folhosas <i>Broadleaves</i>	480.5	(440.6 – 520.3)	99.6
Coníferas <i>Conifers</i>	1.9	(0.0 – 3.9)	0.4
<b>Total</b>	<b>482.4</b>	<b>(442.5 – 522.3)</b>	<b>100.0</b>



20. Peso total da biomassa aérea por grupo de espécies